

THE SM HIGGS-BOSON PRODUCTION IN $\gamma\gamma \rightarrow h \rightarrow b\bar{b}$ AT THE PHOTON COLLIDER AT TESLA

P. NIEŻURAWSKI¹, A. F. ŻARNECKI¹ and M. KRAWCZYK^{2,3}

¹ *Institute of Experimental Physics, Warsaw University, Poland*

² *Theory Division, CERN, Switzerland*

³ *Institute of Theoretical Physics, Warsaw University, Poland*

Abstract

Measuring the $\Gamma(h \rightarrow \gamma\gamma)\text{Br}(h \rightarrow b\bar{b})$ decay at the photon collider at TESLA is studied for a Standard Model Higgs boson of mass $m_h = 120$ GeV. The main background due to the process $\gamma\gamma \rightarrow Q\bar{Q}(g)$, where $Q = b, c$, is estimated using the NLO QCD program (G. Jikia); the results obtained are compared with the LO estimate. Using a realistic luminosity spectrum and performing a detector simulation, we find that $\Gamma(h \rightarrow \gamma\gamma)\text{Br}(h \rightarrow b\bar{b})$ can be measured with an accuracy better than 2% after one year of photon collider running.

A photon collider option of the TESLA e^+e^- collider [1] offers a unique possibility to produce the Higgs boson as an s -channel resonance and to determine its properties with a high accuracy. The neutral Higgs boson couples to the photons through a loop with the massive charged particles, thus $h\gamma\gamma$ coupling is sensitive to contributions of new particles. The SM Higgs boson with a mass below ~ 140 GeV is expected to decay mainly into the $b\bar{b}$ final state. Here we consider the process $\gamma\gamma \rightarrow h \rightarrow b\bar{b}$ for a Higgs-boson mass of $m_h = 120$ GeV at a photon collider at TESLA [2]. Both the signal and background events are generated according to a realistic photon–photon luminosity spectrum. A simulation of the detector response is incorporated as well.

In the analysis we use the CompAZ parametrization [3] of the spectrum [4] to generate energies of the colliding photons. For the energy of primary electrons

$\sqrt{s_{ee}} = 2E_e = 210$ GeV, we obtain a peak of the $J_z = 0$ component of the photon-photon luminosity spectrum at the invariant mass of the two colliding photons $W_{\gamma\gamma}$ equal to 120 GeV. We assume the integrated luminosity of the primary e^-e^- beams equal to $L_{ee}^{geom} = 502 \text{ fb}^{-1}$ [4].

A generation of signal events was done with PYTHIA 6.205 [5], with the parameters for a Higgs boson as in HDECAY [6]. A parton shower algorithm, implemented in PYTHIA, was used to generate the final-state particles. The background events due to processes $\gamma\gamma \rightarrow b\bar{b}(g)$, $c\bar{c}(g)$ were generated using the program written by G. Jikia [7], where a complete NLO QCD calculation for the production of massive quarks is performed within the massive-quark scheme.¹ For a comparison we generated also the LO background events as implemented in PYTHIA. The fragmentation was performed using the PYTHIA program. A fast simulation for a TESLA detector (SIMDET 3.01 [8]) was used to model a detector performance. The jets were reconstructed with the Durham algorithm ($y_{cut} = 0.02$). The double b -tag was required to select the signal $h \rightarrow b\bar{b}$ events ($\varepsilon_{bb} = 70\%$ and $\varepsilon_{cc} = 3.5\%$ were assumed). The following cuts were used to select reconstructed $b\bar{b}$ events: (1) a total visible energy $E_{vis} > 90$ GeV; (2) the ratio of the total longitudinal momentum of all observed particles to the total visible energy $|P_z|/E_{vis} < 0.1$; (3) a number of jets $N_{jets} = 2, 3$; (4) for each jet we require $|\cos \theta_{jet}| < 0.75$. The obtained distributions of the reconstructed $\gamma\gamma$ invariant mass W_{rec} are shown in Fig. 1 (left). The expected relative statistical error of $\Gamma(h \rightarrow \gamma\gamma)\text{Br}(h \rightarrow b\bar{b})$ is equal to $\sqrt{N_{obs}}/(N_{obs} - N_{bkgd})$. If estimated from the selected mass region, it is equal to 1.9%. We introduce the corrected, reconstructed invariant mass as: $W_{corr} \equiv \sqrt{W_{rec}^2 + 2P_T(E_{vis} + P_T)}$. The distributions of the W_{corr} are shown in Fig. 1 (right). In the selected W_{corr} region one achieves an relative accuracy $\Delta \left[\Gamma(h \rightarrow \gamma\gamma)\text{Br}(h \rightarrow b\bar{b}) \right] / \left[\Gamma(h \rightarrow \gamma\gamma)\text{Br}(h \rightarrow b\bar{b}) \right] = 1.7\%$.

Assuming $\text{Br}(h \rightarrow b\bar{b})$ will be measured to 1.5% [9], Higgs-boson partial width $\Gamma(h \rightarrow \gamma\gamma)$ can be extracted with accuracy of 2.3%. Using in addition the result from the e^+e^- Linear Collider for $\text{Br}(h \rightarrow \gamma\gamma)$ [10], one can extract Γ_{tot} with a precision of 10%.

References

- [1] B. Badelek et al., *TESLA TDR*, DESY 2001-011, hep-ex/0108012.
- [2] P. Nieżurawski, A.F. Żarnecki, M. Krawczyk, hep-ph/0208234, accepted by Acta Phys. Polon. B.

¹Other background contributions, from the resolved photon(s) interactions and the overlaying events, were found to be negligible.

- [3] A.F. Żarnecki, hep-ex/0207021.
- [4] V. Telnov, *Nucl. Instrum. Meth.* **A355**, 3 (1995); V. Telnov, talk at the 2nd Workshop of ECFA-DESY Study, Saint Malo, France, April 2002.
- [5] T. Sjöstrand et al., *Comput. Phys. Commun.* **135**, 238 (2001).
- [6] A. Djouadi, J. Kalinowski, M. Spira, *Comput. Phys. Commun.* **108**, 56 (1998).
- [7] G. Jikia, S. Söldner-Rembold, *Nucl. Instrum. Meth.* **A472**, 133 (2001), hep-ex/0101056.
- [8] M. Pohl, H. J. Schreiber, DESY-99-030.
- [9] J-C. Brient, LC-PHSM-2002-003; M. Battaglia, hep-ph/9910271.
- [10] E. Boos et al., *Eur. Phys. J.* **C19**, 455 (2001), hep-ph/0011366.




Figure 1: Reconstructed invariant mass W_{rec} (left) and corrected invariant mass W_{corr} (right) distributions for the selected $b\bar{b}$ events. Contributions of the signal, due to the Higgs boson with a mass $m_h = 120$ GeV, and of the heavy-quark background, calculated in the NLO QCD, are indicated. For comparison, the LO background estimate is also plotted (dots). Arrows indicate the mass window optimized for the measurement of the $\Gamma(h \rightarrow \gamma\gamma)\text{Br}(h \rightarrow b\bar{b})$.